

A GAUSS TYPE FUNCTIONAL EQUATION

SILVIA TOADER, THEMISTOCLES M. RASSIAS, and GHEORGHE TOADER

(Received 30 March 2000)

ABSTRACT. Gauss' functional equation (used in the study of the arithmetic-geometric mean) is generalized by replacing the arithmetic mean and the geometric mean by two arbitrary means.

2000 Mathematics Subject Classification. Primary 39B22; Secondary 26E60.

1. Introduction. By mean we understand a function $M : \mathbb{R}_+ \times \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfies the condition

$$\min(a, b) \leq M(a, b) \leq \max(a, b) \quad \forall a, b > 0. \quad (1.1)$$

The mean is called symmetric if

$$M(a, b) = M(b, a) \quad \forall a, b > 0. \quad (1.2)$$

Usual examples are the power means given by

$$P_n(a, b) = \left(\frac{a^n + b^n}{2} \right)^{1/n} \quad (1.3)$$

for $n \neq 0$, while for $n = 0$ it is the geometric mean

$$P_0(a, b) = G(a, b) = \sqrt{ab}. \quad (1.4)$$

Of course, the arithmetic mean is $A = P_1$.

If M is a mean and $p : \mathbb{R}_+ \rightarrow \mathbb{R}$ is a strictly monotonous function, the expression

$$M(p)(a, b) = p^{-1}[M(p(a), p(b))] \quad (1.5)$$

defines another mean $M(p)$ which is called M -quasi mean (see [1]). For example, the power means are A -quasi means. More exactly $P_n = A(e_n)$, where

$$e_n(x) = x^n \quad \text{for } n \neq 0, \quad e_0(x) = \ln x. \quad (1.6)$$

In what follows, we refer to another famous example of mean. Given two positive numbers a and b , we define successively the terms

$$a_{n+1} = A(a_n, b_n), \quad b_{n+1} = G(a_n, b_n), \quad n \geq 0, \quad (1.7)$$

where $a_0 = a$ and $b_0 = b$. It is known (see [1]) that $(a_n)_{n \geq 0}$ and $(b_n)_{n \geq 0}$ are convergent to a common limit which is denoted by $A \otimes G(a, b)$. It defines the arithmetic-geometric mean of Gauss $A \otimes G$.

The following representation formula is also known (see [1])

$$A \otimes G(a, b) = [f(a, b)]^{-1}, \tag{1.8}$$

where

$$f(a, b) = \frac{1}{2 \cdot \pi} \int_0^{2\pi} \frac{d\theta}{\sqrt{a^2 \cdot \cos^2 \theta + b^2 \cdot \sin^2 \theta}}. \tag{1.9}$$

The proof of this formula is based on the fact that the function f verifies the relation

$$f(A(a, b), G(a, b)) = f(a, b), \tag{1.10}$$

which is called Gauss' functional equation.

These results were generalized as follows. We denote

$$\begin{aligned} r_n(\theta) &= (a^n \cdot \cos^2 \theta + b^n \cdot \sin^2 \theta)^{1/n}, \quad n \neq 0, \\ r_0(\theta) &= \lim_{n \rightarrow 0} r_n(\theta) = a^{\cos^2 \theta} b^{\sin^2 \theta}. \end{aligned} \tag{1.11}$$

If $p : \mathbb{R}_+ \rightarrow \mathbb{R}$ is a strictly monotonous function, then

$$M_{p,n}(a, b) = p^{-1} \left(\frac{1}{2\pi} \int_0^{2\pi} p(r_n(\theta)) d\theta \right) \tag{1.12}$$

defines a symmetric mean. The arithmetic-geometric mean of Gauss is obtained for $n = 2$ and $p(x) = x^{-1}$. For $n = -2$ and $p(x) = x^{-2}$ the mean can be found in [7]. The case $n = 1$ and $p = \log$ was studied in [2]. The essential step was done in [4] by the consideration of the definition (1.12) for $n = 2$ with an arbitrary p . The values $n = -1$ and $n = 1$ were studied in [5, 6]. The general case (of arbitrary n) was studied in [8] and continued in [9]. In [8], Gauss' functional equation was also replaced by a more general equation

$$F(P_q(a, b), P_s(a, b)) = F(a, b). \tag{1.13}$$

In this paper, we generalize the mean (1.12) as well as the functional equation (1.13).

2. An integral mean. We consider the strictly monotonous functions p and q . Using them, we define the functions

$$\begin{aligned} r_q(\theta) &= q^{-1} [q(a) \cdot \cos^2 \theta + q(b) \cdot \sin^2 \theta], \\ f(a, b; p, q) &= \frac{1}{2\pi} \int_0^{2\pi} p[r_q(\theta)] d\theta. \end{aligned} \tag{2.1}$$

It is easy to prove that

$$M_{p,q}(a, b) = p^{-1} [f(a, b; p, q)] \tag{2.2}$$

defines a mean. Choosing $q = e_n$ we obtain $M_{p,q} = M_{p,n}$. We so have generalized the means (1.12). On the other hand, if we let $p \circ q^{-1} = Q$, we have $M_{p,q} = M_{Q,1}(q)$. Thus, $M_{p,q}$ is a $M_{Q,1}$ -quasi mean.

It is thus enough to consider the function

$$f(a, b; p) = \frac{1}{2\pi} \int_0^{2\pi} p(a \cdot \cos^2 \theta + b \cdot \sin^2 \theta) d\theta \tag{2.3}$$

which defines the mean $M_p = M_{p,1}$ by

$$M_p(a, b) = p^{-1}[f(a, b; p)]. \tag{2.4}$$

In what follows, we assume that the function p is two times differentiable. From any of the papers [5, 6, 8, 9], we can deduce the following result.

LEMMA 2.1. *The function f defined by (2.3) has the following partial derivatives:*

$$\begin{aligned} f'_a(c, c; p) &= f'_b(c, c; p) = \frac{1}{2} \cdot p'(c), \\ f''_{aa}(c, c; p) &= f''_{bb}(c, c; p) = \frac{3}{8} \cdot p''(c), \\ f''_{ab}(c, c; p) &= \frac{1}{8} \cdot p''(c). \end{aligned} \tag{2.5}$$

3. The functional equation. We replace (1.13) by a more general functional equation

$$F(M(a, b), N(a, b)) = F(a, b), \tag{3.1}$$

where M and N are two given means. We prove the following result.

LEMMA 3.1. *If the function f , defined by (2.3), verifies the functional equation (3.1), then the function p is a solution of the differential equation*

$$\begin{aligned} p''(c) \cdot \{ [3 \cdot M'_b(c, c) + N'_b(c, c)] \cdot M'_a(c, c) + [M'_b(c, c) + 3 \cdot N'_b(c, c)] \cdot N'_a(c, c) - 1 \} \\ + 4 \cdot p''(c) \cdot [M''_{ab}(c, c) + N''_{ab}(c, c)] = 0. \end{aligned} \tag{3.2}$$

PROOF. Taking in (3.1) the partial derivatives with respect to a , we obtain

$$F'_a[M(a, b), N(a, b)] \cdot M'_a(a, b) + F'_b[M(a, b), N(a, b)] \cdot N'_a(a, b) = F'_a(a, b). \tag{3.3}$$

Taking again the derivatives with respect to b , it follows that

$$\begin{aligned} \{ F''_{aa}[M(a, b), N(a, b)] \cdot M'_b(a, b) + F''_{ab}[M(a, b), N(a, b)] \cdot N'_b(a, b) \} \cdot M'_a(a, b) \\ + \{ F''_{ab}[M(a, b), N(a, b)] \cdot M'_b(a, b) + F''_{bb}[M(a, b), N(a, b)] \cdot N'_b(a, b) \} \cdot N'_a(a, b) \\ + F'_a[M(a, b), N(a, b)] \cdot M''_{ab}(a, b) + F'_b[M(a, b), N(a, b)] \cdot N''_{ab}(a, b) = F''_{ab}(a, b). \end{aligned} \tag{3.4}$$

For $a = b = c$ and the function $F = f$, defined by (2.3), we apply Lemma 2.1 and obtain (3.2). □

CONSEQUENCE 3.2. *If the function f , defined by (2.3), verifies the functional equation (3.1), where the means M and N are symmetric, the function p is a solution of the differential equation*

$$p''(c) + 4 \cdot p'(c) \cdot [M''_{ab}(c, c) + N''_{ab}(c, c)] = 0. \tag{3.5}$$

PROOF. As the means are symmetric, their partial derivatives of the first order are equal to $1/2$ (see [3]), thus (3.2) becomes (3.5). □

CONSEQUENCE 3.3. If the function f , defined by (2.3), verifies the functional equation (1.13), then the function p is given by

$$p(c) = C \cdot c^{r+s-1} + D, \tag{3.6}$$

where C and D are arbitrary constants.

PROOF. We have in (3.5), $M = P_r$ and $N = P_s$. Thus

$$M''_{ab}(c, c) = \frac{1-r}{4 \cdot c}, \quad N''_{ab}(c, c) = \frac{1-s}{4 \cdot c}. \tag{3.7}$$

Replacing them in (3.5), we obtain the differential equation

$$p''(c) + \frac{2-r-s}{c} \cdot p'(c) = 0 \tag{3.8}$$

with the solution given above. □

REMARK 3.4. This last result was obtained in [8]. As it is shown in [9], the condition is also sufficient for $r = -s = 1$.

REMARK 3.5. Equation (3.1) can be further generalized at

$$F(g(M(a, b)), g(N(a, b))) = h(F(a, b)), \tag{3.9}$$

where g and h are two given functions. We have in view the following result given in [2]. The function f , defined by (2.3), verifies the relation

$$f(A^2(a, b), G^2(a, b); \log) = 2 \cdot f(a, b; \log). \tag{3.10}$$

4. Special means. A problem which is studied for the integral means defined in [4, 5, 6, 8, 9] is that of the determination of the cases in which the mean reduces at a given one, usually a power mean. Similar results can be given also in more general circumstances. We prove the following lemma.

LEMMA 4.1. *If for a given mean N , we have $M_p = N$, then the function p verifies the equation*

$$p''(c) \cdot [8 \cdot N'_a(c, c) \cdot N'_b(c, c) - 1] + 8 \cdot p'(c) \cdot N''_{ab}(c, c) = 0. \tag{4.1}$$

PROOF. In the given hypotheses, we have

$$f(a, b; p) = p[N(a, b)]. \tag{4.2}$$

Taking the partial derivatives with respect to a , we have

$$f'_a(a, b; p) = p'[N(a, b)] \cdot N'_a(a, b). \tag{4.3}$$

Then we take the derivative with respect to b , we obtain

$$f''_{ab}(a, b; p) = p''[N(a, b)] \cdot N'_a(a, b) \cdot N'_b(a, b) + p'[N(a, b)] \cdot N''_{ab}(a, b). \tag{4.4}$$

For $a = b = c$, Lemma 2.1 gives (4.1). □

CONSEQUENCE 4.2. If we have $M_p = N$, with the symmetric mean N , then the function p verifies the equation

$$p''(c) + 8 \cdot p'(c) \cdot N_{ab}''(c, c) = 0. \quad (4.5)$$

CONSEQUENCE 4.3. If we have $M_p = P_r$, then the function p is given by

$$p(c) = C \cdot c^{2r-1} + D, \quad (4.6)$$

where C and D are arbitrary constants.

REMARK 4.4. In [9], it is shown that the above condition is also sufficient for $r = 0$, $1/2$, and 1 .

REFERENCES

- [1] P. S. Bullen, D. S. Mitrinović, and P. M. Vasić, *Means and their Inequalities*, Mathematics and its Applications (East European Series), vol. 31, D. Reidel Publishing Co., Massachusetts, 1988, translated and revised from the Serbo-Croatian. [MR 89d:26003](#). [Zbl 687.26005](#).
- [2] B. C. Carlson, *Invariance of an integral average of a logarithm*, Amer. Math. Monthly **82** (1975), 379–382. [MR 50#13628](#). [Zbl 301.33001](#).
- [3] D. M. E. Foster and G. M. Phillips, *A generalization of the Archimedean double sequence*, J. Math. Anal. Appl. **101** (1984), no. 2, 575–581. [MR 85g:40001](#). [Zbl 555.40002](#).
- [4] H. Haruki, *New characterizations of the arithmetic-geometric mean of Gauss and other well-known mean values*, Publ. Math. Debrecen **38** (1991), no. 3-4, 323–332. [MR 92f:39014](#). [Zbl 746.39004](#).
- [5] H. Haruki and T. M. Rassias, *A new analogue of Gauss' functional equation*, Int. J. Math. Math. Sci. **18** (1995), no. 4, 749–756. [MR 97a:39025](#). [Zbl 855.39019](#).
- [6] ———, *New characterizations of some mean-values*, J. Math. Anal. Appl. **202** (1996), no. 1, 333–348. [MR 97f:39026](#). [Zbl 878.39005](#).
- [7] G. Pólya and G. Szegő, *Isoperimetric Inequalities in Mathematical Physics*, Annals of Mathematics Studies, no. 27, Princeton University Press, New Jersey, 1951. [MR 13,270d](#). [Zbl 044.38301](#).
- [8] G. Toader, *Some mean values related to the arithmetic-geometric mean*, J. Math. Anal. Appl. **218** (1998), no. 2, 358–368. [MR 98m:26017](#). [Zbl 892.26015](#).
- [9] G. Toader and T. M. Rassias, *New properties of some mean values*, J. Math. Anal. Appl. **232** (1999), no. 2, 376–383. [MR 2000c:26020](#). [Zbl 924.39016](#).

SILVIA TOADER: DEPARTMENT OF MATHEMATICS, TECHNICAL UNIVERSITY, RO-3400 CLUJ, ROMANIA

THEMISTOCLES M. RASSIAS: DEPARTMENT OF MATHEMATICS, NATIONAL TECHNICAL UNIVERSITY, 15780 ATHENS, GREECE
E-mail address: trassias@math.ntua.gr

GHEORGHE TOADER: DEPARTMENT OF MATHEMATICS, TECHNICAL UNIVERSITY, RO-3400 CLUJ, ROMANIA
E-mail address: gheorghe.toader@math.utcluj.ro

Special Issue on Time-Dependent Billiards

Call for Papers

This subject has been extensively studied in the past years for one-, two-, and three-dimensional space. Additionally, such dynamical systems can exhibit a very important and still unexplained phenomenon, called as the Fermi acceleration phenomenon. Basically, the phenomenon of Fermi acceleration (FA) is a process in which a classical particle can acquire unbounded energy from collisions with a heavy moving wall. This phenomenon was originally proposed by Enrico Fermi in 1949 as a possible explanation of the origin of the large energies of the cosmic particles. His original model was then modified and considered under different approaches and using many versions. Moreover, applications of FA have been of a large broad interest in many different fields of science including plasma physics, astrophysics, atomic physics, optics, and time-dependent billiard problems and they are useful for controlling chaos in Engineering and dynamical systems exhibiting chaos (both conservative and dissipative chaos).

We intend to publish in this special issue papers reporting research on time-dependent billiards. The topic includes both conservative and dissipative dynamics. Papers discussing dynamical properties, statistical and mathematical results, stability investigation of the phase space structure, the phenomenon of Fermi acceleration, conditions for having suppression of Fermi acceleration, and computational and numerical methods for exploring these structures and applications are welcome.

To be acceptable for publication in the special issue of Mathematical Problems in Engineering, papers must make significant, original, and correct contributions to one or more of the topics above mentioned. Mathematical papers regarding the topics above are also welcome.

Authors should follow the Mathematical Problems in Engineering manuscript format described at <http://www.hindawi.com/journals/mpe/>. Prospective authors should submit an electronic copy of their complete manuscript through the journal Manuscript Tracking System at <http://mts.hindawi.com/> according to the following timetable:

Manuscript Due	December 1, 2008
First Round of Reviews	March 1, 2009
Publication Date	June 1, 2009

Guest Editors

Edson Denis Leonel, Departamento de Estatística, Matemática Aplicada e Computação, Instituto de Geociências e Ciências Exatas, Universidade Estadual Paulista, Avenida 24A, 1515 Bela Vista, 13506-700 Rio Claro, SP, Brazil ; edleonel@rc.unesp.br

Alexander Loskutov, Physics Faculty, Moscow State University, Vorob'evy Gory, Moscow 119992, Russia; loskutov@chaos.phys.msu.ru